

DOI: https://doi.org/10.1093/eep/dvae018 Advance Access Publication 23 September 2024 Research Article

DNA methylation correlates with transcriptional noise in response to elevated pCO₂ in the eastern oyster (Crassostrea virginica)

Yaamini R. Venkataraman^{1,*}, Ariana S. Huffmyer^{2,3}, Samuel J. White², Alan Downey-Wall⁴, Jill Ashey³, Danielle M. Becker³, Zachary Bengtsson², Hollie M. Putnam³, Emma Strand^{3,5}, Javier A. Rodríguez-Casariego⁶, Shelly A. Wanamaker⁵, Katie E. Lotterhos⁷, Steven B. Roberts¹

¹Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, United States

- ²School of Aquatic and Fisheries Sciences, University of Washington, Seattle, WA 98195, United States
- ³Department of Biological Sciences, University of Rhode Island, Kingston, RI 02881, United States

⁴Embark Veterinary, Boston, MA 02111, United States

⁵Gloucester Marine Genomics Institute, Gloucester, MA 01930, United States

⁶Environmental Epigenetics Laboratory, Institute of Environment, Florida International University, Miami, FL, 33199, United States

⁷Northeastern University Marine Science Center, Nahant, MA 01908, United States

*Corresponding author. Biology Department, Woods Hole Oceanographic Institution, 266 Woods Hole Rd MS #33, Woods Hole, MA 02543, United States. E-mail: yaamini.venkataraman@whoi.edu

Abstract

Ocean acidification significantly affects marine calcifiers like oysters, warranting the study of molecular mechanisms like DNA methylation that contribute to adaptive plasticity in response to environmental change. However, a consensus has not been reached on the extent to which methylation modules gene expression, and in turn plasticity, in marine invertebrates. In this study, we investigated the impact of pCO_2 on gene expression and DNA methylation in the eastern oyster, *Crassostrea virginica*. After a 30-day exposure to control (572 ppm) or elevated pCO_2 (2827 ppm), whole-genome bisulfite sequencing (WGBS) and RNA-seq data were generated from adult female gonad tissue and male sperm samples. Although differentially methylated loci (DMLs) were identified in females (89) and males (2916), there were no differentially expressed genes and only one differentially expressed transcript in females. However, gene body methylation impacted other forms of gene activity in sperm, such as the maximum number of transcripts expressed per gene and changes in the predominant transcript expressed. Elevated pCO_2 exposure increased gene expression regulation. Functional annotation of genes with changes in transcript-level expression or containing DMLs revealed several enriched biological processes potentially involved in elevated pCO_2 response, including apoptotic pathways and signal transduction, as well as reproductive functions. Taken together, these results suggest that DNA methylation may regulate gene expression variability to maintain homeostasis in elevated pCO_2 conditions and could play a key role in environmental resilience in marine invertebrates.

Keywords: DNA methylation; gene expression; oysters; ocean acidification; reproduction

Introduction

Ocean acidification (OA) is a pressing environmental issue resulting from the increased absorption of atmospheric carbon dioxide (CO_2) by the world's oceans [1]. Increased concentrations of oceanic CO_2 lead to decreases in seawater pH, lower calcium carbonate concentrations, and reductions in aragonite saturation state [1]. These environmental alterations pose a significant threat to marine ecosystems, particularly to calcifying organisms such as oysters [2–5]. Reductions in saturation state can compromise calcification and increase the energetic cost of building shells [6–9] and can impact broad physiological pathways such as protein synthesis, energy production, metabolism, antioxidant responses, and reproduction [10–14]. Since oysters are vital for biodiversity, aquaculture, and coastal economies [15–17], understanding the biological responses of these organisms to OA is critical for assessing ecological impacts and developing strategies to mitigate negative effects of OA in oyster restoration and aquaculture production.

The ability of organisms to respond to and cope with OA is dependent on complex changes in whole-organism physiology,

Received 4 April 2024; revised 15 August 2024; accepted 22 September 2024 © The Author(s) 2024. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (https://creativecommons.org/ licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site–for further information please contact journals.permissions@oup.com. behavior, and molecular mechanisms [4, 18-21]. Additionally, plasticity, or the capacity for an organism to adjust its physiology to maintain function and performance in response to environmental change [22], is another important factor governing responses to OA. Plastic responses to environmental stressors can also be inherited, providing a mechanism for the parent to pass on "environmental memory," through mechanisms such as alterations in reproduction and offspring provisioning [23]. There are several examples of adult oyster exposure to OA yielding beneficial plasticity in offspring. For example, Sydney rock oysters (Saccostrea glomerata) exposed to high pCO₂ during reproductive conditioning yielded larger and faster-developing larvae compared to counterparts in ambient conditions [24, 25], Olympia oyster (Ostrea lurida) offspring with parental history of OA exposure had better performance than ambient counterparts when outplanted [26], and exposure of adult eastern oysters (Crassostrea virginica) to elevated pCO₂ increased larval growth rates [27]. However, parental exposure to OA in the Pacific oyster (Crassostrea gigas), northern quahog (Mercenaria mercenaria), and bay scallop (Argopecten irradians) led to poorer larval survival [28, 29]. These contrasting results demonstrate that the capacity for intergenerational plasticity is complex and requires additional research. As adult reproductive tissue gives rise to gametes and offspring, it is critical to examine the molecular mechanisms impacting reproductive homeostasis in response to OA.

Epigenetic modifications are a potential mediator of transgenerational plastic responses to environmental change. These modifications alter regulation of gene expression without altering the DNA sequence itself and include multiple interacting mechanisms such as DNA methylation, chromatin remodeling, histone modifications, and noncoding RNAs [30]. DNA methylation-the addition of a methyl group to the cytosine adjacent to a guanine in a CpG dinucleotide [30, 31]—is the most studied epigenetic mechanism in marine invertebrates [32]. DNA methylation can change expression of genes by altering the binding of transcriptional machinery and can affect variability of transcription [31]. Changes in methylation are thought to be associated with phenotypic changes, with several examples of this connection illustrated in mammalian literature [33]. Several studies in marine invertebrates have also demonstrated the potential for methylomes (i.e. information on DNA methylation in a genome) to be inherited by offspring [34–38]. Methylomes are responsive to OA across a wide variety of taxa, such as pteropods [39], copepods [36, 37], corals [40], sea urchins [41], and oysters [42–46]. These methylation responses are often species specific, even within the same taxa [33]. For example, genes involved in protein ubiquitination, cytoskeletal regulation, and metabolism were all differentially methylated in various tissues and species, including the mantle [42] and gonad [44] of C. virginica, the mantle [45] and larvae [43, 47] of the Hong kong oyster (Crassostrea hongkongensis), and the reproductive tissue [46] of female C. gigas. Genes with differential methylation were not necessarily orthologous across species. However, common processes impacted by DNA methylation across species and tissue types suggest a role for methylation in molluscan OA response.

Even though methylation is environmentally responsive, direct connections between methylation and plasticity are not readily apparent in molluscs. Where gene expression has been used as a proxy for plasticity, molluscan studies report no relationship between differential gene expression and genes with differential methylation in response to environmental stress [34, 42, 45, 48, 49]. One exception in molluscan literature is Dang et al. [47], which found significant overlap between differentially expressed genes (DEGs) and differentially methylated genes in juvenile C. hongkongensis after exposure to OA conditions as larvae. Differentially expressed and methylated genes were often methylated in introns and had roles in signal transduction [47]. Studies that do not explore gene expression and methylation have found associations between methylation and phenotype, although causal relationships have not been established. While differences in mantle methylomes cannot be linked to changes in calcification [42, 45], there is evidence associating differential methylation with the ability to reproduce in stressful OA conditions [46], which in turn could be linked to maladaptive plasticity in larval C. gigas [29]. Additionally, differential methylation of C. hongkongensis larvae exposed to elevated pCO₂ was associated with higher metamorphosis rates but poor substratum selection [43]. Taken together, this suggests that environmentally responsive DNA methylation may play a role in modulating phenotypic plasticity in adult reproductive tissue and could contribute to intergenerational carryover effects. Therefore, it is important to investigate knowledge gaps surrounding the role of methylation in altering gene expression and plasticity in oysters exposed to OA during reproductive conditioning.

This study explored the impact of OA on gene expression and DNA methylation in C. virginica adult reproductive tissue and the relationship between gene expression and methylation (Fig. 1). We used oysters from an accompanying study, which found that parental exposure to elevated pCO₂ increased growth rates in larvae, especially when those larvae were also reared in elevated pCO₂ conditions [27]. Female oysters in control and elevated pCO₂ conditions did not have any differences in egg quality or size, suggesting that epigenetic mechanisms may have mediated the observed carryover effect [27]. We build on previous work demonstrating environmentally sensitive methylation in Crassostrea spp. reproductive tissue [44, 46] by examining sex-specific effects of OA on reproductive tissue responses. As female and male oysters undergo distinct gametogenic processes, and egg and sperm have unique roles in reproduction, it follows that there are sex-specific methylation [50-52] and gene expression [53] patterns in reproductive tissue that would confound findings if analyzed together. We compared female gonad tissue and male sperm exposed to control and elevated pCO₂ conditions to investigate (i) the influence of genetic variation on methylation and gene expression; (ii) the impact of OA on changes in gene activity and methylation; and (iii) the functional role of DNA methylation in regulating gene activity. We define gene activity as encompassing genelevel expression, but also transcript-level processes such as transcript expression, changes in the number of unique transcripts expressed per gene, shifts in the predominant isoform expressed, and alternative splicing. By incorporating transcript-level analyses, we can further elucidate the functional role of methylation in regulating gene expression, and therefore plasticity, in response to environmental change. Through this comprehensive approach, we seek to uncover the molecular basis of oyster responses to OA, providing insights into potential impacts on reproduction and the molecular mechanisms involved in marine invertebrate stress response.



Figure 1. Conceptual diagram of study design.

Results

Seawater chemistry

As previously reported by McNally *et al.* [27], seawater of control and elevated pCO_2 conditions were 572 and 2827 ppm, respectively. The $\Omega_{aragonite}$ of seawater differed significantly between control and elevated pCO_2 conditions (Welch's two-sample t-test: t = 12.33, df = 7, P-value < .0001).

Genetic variation

The full whole-genome bisulfite sequencing (WGBS) dataset of all samples (female_{control} = 8, female_{exposed} = 8, male_{control} = 4, $male_{exposed} = 6$) was used to examine genetic variation and the influence of that variation on gene methylation and expression. Even though the full dataset contains different cell types (i.e. female gonad tissue and male sperm), we wanted to determine the relatedness between female and male oysters. After filtering, we kept 2343637 single nucleotide polymorphisms (SNPs) out of 144873997 possible sites to generate a pairwise relatedness matrix and calculate genetic distance. Pairwise genetic, methylation, and gene expression distances were calculated using Euclidean distances. There was no significant correlation between pairwise genetic distance and pairwise methylation distance (R = -0.16, P-value = .99) (Supplementary File S1) nor genetic distance and pairwise gene expression distance (R=-0.053, Pvalue = .76) (Supplementary File S1). Because pairwise genetic distance between samples was not related to pairwise distance in methylation nor gene expression between samples, genetic distance was not incorporated in downstream analyses of methylation and gene expression.

Gene activity

Gene and transcript expression

Across all female and male samples, 67.57% of RNA reads aligned to the C. *virginica* genome. DEGs and differentially expressed transcripts (DETs) were identified between control and OA-exposed samples in female gonad tissue and male sperm samples separately. There were no DEGs in females, but a single DET was identified (Supplementary File S2). The transcript was associated with gene ID LOC111134855 (uncharacterized). No DEGs nor DETs were found in males.

Changes in the maximum number of transcripts expressed per gene between control and OA-exposed samples were characterized as another measure of gene activity. In female gonads, a total of 3966 genes (~10% of total expressed genes) showed differences in the maximum number of transcripts expressed between female control and OA-exposed samples (Table 1). Under OA conditions in female gonads, 1529 (38.6%) genes had a greater number of transcripts expressed per gene, while 2437 had fewer transcripts expressed per gene (Table 1; Supplementary File S3). A total of 211 biological processes were enriched in genes with more transcripts under elevated pCO₂, with the top 10 significantly enriched gene ontology (GO) terms including establishment of localization (Pvalue = 2.0×10^{-10}) and cell adhesion (P-value = 2.6×10^{-10}). Genes with less transcripts expressed under elevated pCO₂ conditions had 469 enriched biological processes. The top 10 significantly enriched GO terms included anatomical structure development $(P-value = 5.3 \times 10^{-14})$ and signaling $(P-value = 7.0 \times 10^{-13})$. Statistical output for all GO terms can be found in Supplementary File S4.

In sperm, 3123 (72.2%) of genes that exhibited changes in the maximum number of transcripts had a greater number of transcripts expressed between control and OA-exposed samples, while 1204 (27.8%) had fewer transcripts expressed in elevated pCO₂ conditions (Table 1; Supplementary File S3). Genes with more transcripts expressed in elevated pCO₂ conditions in sperm had 391 over-represented biological processes, with the top 10 significantly enriched GO terms including signaling (P-value = 4.9×10^{-9}) and cytoskeletal organization (P-value = 1.9×10^{-8}) (Supplementary File S4). Genes with fewer transcripts expressed in elevated pCO₂ conditions had 238 enriched biological processes, with the

Table 1. Gene and transcript expression patterns attributable to elevated pCO₂ conditions

Sex	Total genes expressed	Genes with more transcripts expressed	Genes with less transcripts expressed	Genes with ≥ 2 transcripts	Genes with a shift in predominant transcript	
Female	39 047	1529	2437	38 575	2252	
Male	39 299	3123	1204	10 559	1768	

Changes in the maximum number of transcripts are described with respect to elevated pCO_2 exposure, with increases in the maximum number of transcripts representing more transcripts expressed in elevated pCO_2 -exposed samples and decreases representing more transcripts expressed in control samples. Genes with multiple (≥ 2) transcripts were considered for predominant transcript analyses.

top 10 significantly enriched GO terms including organelle organization (P-value = 4.0×10^{-7}) and intracellular signal transduction (P-value = 1.6×10^{-6}) (Supplementary File S4).

Predominant transcript identification

Low pH-induced shifts in the predominant transcript expression were assessed separately for female reproductive tissue and male sperm, where the predominant transcript was defined as the most highly expressed transcript within a gene. Of the 38575 genes that had expression data for multiple transcripts in female gonads, 2252 genes (5.83%) had shifts in the predominant transcript (Table 1; Supplementary File S3). A total of 226 biological process GO terms were over-represented in genes where the predominant transcript shifted, with the top 10 significantly enriched GO terms including regulation of small GTPase-mediated signaling (P-value = 6.2×10^{-9}), regulation of signal transduction (P-value = 5×10^{-6}), and reproduction (P-value = 8.1×10^{-6}). Statistical output for all GO terms can be found in Supplementary File S4.

For sperm, 1768 (16.7%) out of 10559 genes with data for multiple transcripts had shifts in the predominant transcript (Table 1; Supplementary File S3). Thirty-three biological processes were enriched in genes with shifts in the predominant transcript. The top 10 significantly enriched terms included actomyosin structure organization (P-value = .0015) and chemotaxis (P-value = .0017) (Supplementary File S4).

Alternative splicing

For each sex, we described patterns of alternative splicing by testing for variation in expression across exon position and treatment using ANOVA-simultaneous component analyses (ASCAs) [54], calculated as the expression of exons 2-6 relative to expression of the first exon for every gene in each sample. ASCAs identified groups of genes with shared expression patterns across exons (e.g. genes that have highest expression of exon 1 relative to other exons and genes that have highest expression of exons 1 and 2 relative to other exons). Principal components (PCs) of the ASCA model describe variation in patterns of gene expression across exons. Of the 10 PCs identified by the ASCA model for each sex, PCs 1-6 explained >1% of the variation in relative exon expression and were selected for further analysis (Supplementary File S5). In both females (Supplementary File S6) and males (Supplementary File S7), PCs 1–4 described alternative splicing patterns that did not differ by treatment and explained the majority of variation in alternative splicing patterns (91.77% in females, 86.72% in males) (Supplementary File S5). In contrast, PCs 5 and 6 described alternative splicing patterns that differed between treatments in females (Fig. 2) and males (Fig. 3). Patterns of alternative splicing described by PCs 5 and 6 accounted for low variance relative to PCs 1-4 (7.69% in females, 10.89% in males; Supplementary File S5). Due to the low variance explained by PC 6 specifically (\sim 1.5%), confidence intervals could not be calculated for this pattern.

There were four distinct patterns of alternative splicing that were not affected by treatment and were consistent between females (Supplementary File S6) and males (Supplementary File S7). Specifically, PC 1 described genes with expression that changes linearly across exon position, which includes genes that have both increasing and decreasing expression across exon position (females: Supplementary File S6AB, males: Supplementary File S7AB). PC 2 described genes that have either increased or decreased expression in middle exons (females: Supplementary File S6CD, males: Supplementary File S7CD), and PCs 3 and 4 described genes that have either higher or lower expression in other combinations of exon positions (females: Supplementary File S6E-H, males: Supplementary File S7E-H). Since these patterns did not differ between treatments, we did not conduct functional enrichment analyses and proceeded to analyze alternative splicing patterns that showed treatment differences (i.e. PCs 5 and 6).

There were two patterns of alternative splicing that revealed differences between treatments and were consistent in both females (Fig. 2) and males (Fig. 3). PC 5 described genes that show differential expression of exons 2–6 relative to exon 1 between treatments (Figs 2a and 3a). We visualized the top 10 genes that were most strongly associated with this pattern, revealing genes that show either lower or higher expression in exons 2–6 in the exposed treatment (Figs 2b and c and 3b and c). PC 6 described genes that show differential expression of front and/or end exons relative to exon 1 (Figs 2d and 3d). We also visualized the top 10 genes associated with this pattern in females (Fig. 2f) and males (Fig. 3f), which revealed genes that had differential expression in exons 2–4 or exons 5 and 6 between treatments. Although the patterns were consistent across sexes, the top genes associated with these patterns differed.

We further analyzed the top 20 genes associated with PCs 5 and 6. For females, five of the top 20 genes from PCs 5 and 6 had changes in the maximum number of transcripts expressed (Supplementary File S3). There were 38 enriched biological process GO terms identified in the top female genes from PCs 5 and 6. Statistical output for all GO terms can be found in Supplementary File S4. The top 10 significantly enriched processes included protein UFMylation (ubiquitin-like modification; P-value = 6.1×10^{-5}) and manganese ion transmembrane transport (P-value = .003) (Supplementary File S4). In males, four of the top 20 genes from PCs 5 and 6 had changes in the maximum number of transcripts in males (Supplementary File S3). A total of 26 over-represented biological processes were identified, with the top 10 significantly enriched GO terms including regulation of metallopeptidase activity (P-value = .004) and acetyl-CoA metabolic process (P-value = .003).



Figure 2. PCs identified by ASCA describing patterns of alternative splicing that show differential effects of treatment in females. (a) PC 5 characterizes genes with differential expression in exons 2–6 relative to exon 1 between treatments. (b) Top 10 genes associated with alternative splicing patterns characterized by PC 5, ordered by PC loading score, with error bars indicating 95% confidence intervals with variables indicating genes. (c) Expression (z-score) of exon position for each of the top 10 genes associated with PC 5. (d) PC 6 characterizes genes with differential expression in exons 2–4 relative to exon 1 or in exons 5 and 6 relative to exon 1 between treatments (confidence intervals could not be calculated for PC 6 and are therefore not shown). (e) Top 10 genes associated with alternative splicing patterns characterized by PC 6, ordered by PC loading score. (f) Expression (z-score) of exon position for each of the top 10 genes associated with PC 6.

DNA methylation analysis

Approximately 57% of the quality trimmed reads were aligned to the genome, and on average 12.4% of the CpG loci were methylated. The average number of loci with a minimum of 10× coverage in 26 individuals was 8.1M (56% of 14458703 total CpGs in the C. virginica genome). The baseline methylation landscape was characterized for each sex separately. A total of 12678572 and 11991497 non-C \rightarrow T SNP CpGs had at least 10× coverage in at least one female or male sample, respectively. For both sexes, a majority of CpGs were lowly methylated (females: 10212421 CpGs, 80.5%; males: 9657277 CpGs, 80.5%), followed by highly methylated (females: 1254583 CpGs, 9.9%; males: 1467485 CpGs, 12.2%) and moderately methylated (females: 1211567 CpGs, 9.6%; males: 866734 CpGs, 7.2%) (Supplementary File S8).

In females, highly methylated CpGs were found primarily in genes, with 662 082 CpGs (52.8%) in introns, 416 646 CpGs in coding sequences (CDS) (33.2%), and 62 784 CpGs in untranslated regions of exons (exon UTR) (5.0%) (Supplementary File S8). More highly methylated CpGs were located in the 1000 kb flanks downstream of genes (40 697 CpGs; 3.2%) than in flanks upstream of transcription start sites (7896 CpGs; 0.6%). A total of 75 013 CpGs (6.0%) were found in intergenic regions, 80 887 CpGs (6.4%) in transposable elements (TE), and 21 399 CpGs (1.7%) in lncRNA. Similarly, highly methylated CpGs in males were concentrated in genes, with more CpGs located in introns (812 892 CpGs, 55.4%) than CDS (438 908 CpGs, 30.0%) or exon UTR (77 872 CpGs, 5.3%) (Supplementary File S8). Downstream flanks contained more CpGs (46 650 CpGs, 3.2%) than upstream flanks (10 088 CpGs, 0.7%). Males also

had higher concentrations of CpGs in TE (93 876 CpGs, 6.4%), followed by intergenic regions (92 758 CpGs, 6.3%) and lncRNA (27 706 CpGs, 1.9%). The distribution of highly methylated CpGs in both sexes in genome features was significantly different than the distribution of all CpGs in the genome (chi-squared contingency tests; females: Supplementary File S9, males: Supplementary File S10).

Differential methylation at the CpG locus-level

There were 128 putative differentially methylated loci (DMLs) in female gonads and 4175 putative DMLs in male sperm when comparing elevated pCO₂-exposed and control oysters. After removing $C \rightarrow T$ SNPs, there were 89 and 2916 female and male DMLs, respectively. Female DMLs were found predominantly in genes (74 DMLs; 83.1%), with 37 DMLs in introns (41.6%), 29 DMLs in CDS (32.6%), and 8 DMLs in exon UTR (9.0%) (Fig. 4). Seven DMLs (7.9%) were found in intergenic regions, four DMLs (4.5%) were found in both upstream and downstream flanks, and one DML (1.1%) was found in both lncRNA and TE. The proportion of female DMLs found in either CDS or intergenic regions was significantly different from the proportion of 10× CpGs in those same genome features (chi-squared contingency test results can be found in Supplementary File S11). GO enrichment analysis revealed 85 biological process GO terms enriched in genes containing DMLs. The top 10 significantly enriched processes included dolichol metabolic process and (P-value = 1.1×10^{-5}) negative regulation of apoptotic process (P-value = .001). Statistical output for all GO terms can be found in Supplementary File S4.



Figure 3. PCs identified by ASCA describing patterns of alternative splicing that show differential effects of treatment in males. (a) PC 5 characterizes genes with differential expression in exons 2–6 relative to exon 1 between treatments. (b) Top 10 genes associated with alternative splicing patterns characterized by PC 5, ordered by PC loading score, with error bars indicate 95% confidence intervals with variables indicating genes. (c) Expression (z-score) of exon position for each of the top 10 genes associated with PC 5. (d) PC 6 characterizes genes with differential expression in the exons 2–4 relative to exon 1 or in exons 5 and 6 relative to exon 1 between treatments (confidence intervals could not be calculated for PC 6 and are therefore not shown). (e) Top 10 genes associated with PC 6.

Similar to females, the majority of male DMLs were found in gene bodies (2322 DMLs; 79.6%) (Fig. 4). Within genes, 1332 DMLs were found in introns (45.7%), 815 DMLs in CDS (27.9%), and 190 DMLs (6.5%) in exon UTR. More DMLs were found in downstream flanks (163 DMLs; 5.6%) than upstream flanks (60 DMLs; 2.1%). A total of 279 DMLs (9.6%) were found in intergenic regions, while 144 DMLs (4.9%) were in TE and 52 DMLs (1.8%) were in lncRNA. The proportion of male DMLs in exon UTR, CDS, introns, upstream flanks, downstream flanks, and intergenic regions differed significantly from the proportion of 10× CpGs in those genome features (chi-squared contingency test results can be found in Supplementary File S12). A total of 359 biological process GO terms were enriched in genes containing DMLs, with the top 10 over-represented processes including organelle organization (P-value = 7.1×10^{-15}), regulation of GTPase activity (P-value = 1.2×10^{-11}), and macromolecule localization (Pvalue = 9.8×10^{-9}) (Supplementary File S4).

Methylation influence on gene activity

A significant and positive correlation was found between gene methylation and expression distance matrices, indicating that samples that were highly similar in gene expression were also highly similar in methylation (R=0.843, P-value=.01) (Supplementary File S1). To investigate the reason for this higher-level correlation, methylation's influence on gene activity was evaluated by examining its effect on (i) gene and transcript expression, (ii) predominant transcript shifts, (iii) transcriptional noise, and (iv) alternative splicing.

Influence of methylation on gene and transcript expression

The one transcript differentially expressed between control and elevated pCO₂-exposed female oysters did not contain DMLs. Since a one-to-one relationship was not identified between DML and DET, the influence of methylation on other aspects of gene activity was explored. We first examined connections between the average gene methylation and changes in the maximum number of transcripts expressed. The number of DMLs in genes with changes in the maximum number of expressed transcripts ranged from zero to two in females (Supplementary File S13) and 0 to 25 in males (Supplementary File S13). A binomial generalized linear model (GLM) was used to evaluate the influence of the explanatory variable, gene body methylation, on whether or not there was a change in the response variable, the maximum number of transcripts expressed due to elevated pCO₂ exposure. The number of DMLs in a gene, gene length, and gene expression were used as additional explanatory variables. The number of DMLs in a gene was not a significant predictor of a change in maximum transcripts expressed (Table 2; Supplementary File S14) in female gonad tissue or male sperm. Changes in gene body methylation had a marginally significant negative effect on changes in the maximum number of transcripts male oysters (Table 2). For both sexes, changes in gene expression and gene length were significant predictors (Table 2).

Influence of methylation on predominant transcript shifts

Several DMLs were found in genes with a shift in the predominant transcript expressed. In females, zero to three DMLs were found



Figure 4. Genomic locations of 89 DMLs in female gonad tissue and 2916 DMLs in male sperm. (a) Proportion 10× CpGs with data versus DMLs in various genome features. (b) Total number of DMLs in various genome features.

Table 2. Summary of binomial GLMs for female and male oysters examining the influence of the explanatory variables of gene body methylation, DML count, gene length, and gene expression on the response variable of changes to the maximum number of transcripts expressed due to elevated pCO_2 exposure

	Females			Males		
Explanatory variable	eta Estimate	z-value	P-value	eta Estimate	z-value	P-value
Intercept	0.24	0.67	0.51	2.66	6.88	6.00 × 10 ⁻¹² **
Change in gene body methylation	0.007	0.44	0.66	-0.01	-1.76	0.08
DML count	0.35	0.23	0.82	-0.05	-1.07	0.29
log ₂ (Change in gene expression)	0.65	11.31	1.13 × 10 ^{-29**}	-0.54	-5.86	4.63 × 10 ^{-9**}
log ₁₀ (Gene length)	-0.19	-2.14	0.03**	0.18	6.27	3.67 × 10 ^{-10**}

*P-value < .01. **P-value < .05.

in these genes (Supplementary File S13), with 0–25 DMLs in males (Supplementary File S13). Binomial GLMs were used to quantify the impact of gene body methylation, DML count, gene expression, and gene length on whether or not there was a shift in the predominant transcript expressed due to low pH exposure. In female oysters, change in gene body methylation and number of DMLs in genes did not have a significant effect (Table 3; Supplementary File S15). However, change in gene body methylation had a significant positive effect on a shift in the predominant transcript in males (Table 3; Supplementary File S15). Gene expression and gene length were significant predictors of predominant transcript shift in both females and males (Table 3).

Transcriptional noise analysis

Models demonstrate that both gene body methylation and expression have a significant influence on gene expression variability, or transcriptional noise (Table 4; Fig. 5). All predictors were included in the best-fit model for each sex. For female gonads (model adjusted $R^2 = 0.60$), transcriptional noise was negatively correlated with treatment (P-value = .002; Fig. 5a), gene expression (P-value = 0; Fig. 5c), gene body methylation (P-value = 0; Fig. 5e), and the interaction between treatment and expression, but positively correlated with methylation variation (P-value = 1.87 × 10⁻²⁷²; Fig. 5g), gene expression squared, the interaction between treatment and gene expression squared, and

Table 3. Summary of binomial GLMs for female and male oysters examining the influence of the explanatory variables of gene body methylation, DML count, gene length, and gene expression on the response variable, a shift in the predominant transcript due to elevated pCO₂ exposure

Explanatory variable	Females β Estimate	z-value	P-value	Males eta Estimate	z-value	P-value
Intercept	-8.49	-35.93	9.54 × 10 ^{-283**}	-3.04	-9.44	3.71×10 ^{-21**}
Change in gene body methylation	-0.005	-0.58	0.56	0.01	2.03	0.04
DML count	-11.64	-0.10	0.92	0.07	1.61	0.11
log ₂ (Change in gene expression) log ₁₀ (Gene length)	-0.21 1.45	-11.10 25.18	$7.20 \times 10^{-140^{**}}$ $1.32 \times 10^{-28^{**}}$	0.05 0.33	2.16 4.28	0.03 ^{**} 1.88 × 10 ^{-5**}

*P-value < .01. **P-value < .05.

gene length (Table 4). Inclusion of polynomial terms for gene expression reflects that gene expression variability is the highest at very low and very high levels of gene expression. Methylation had the highest partial R^2 (partial $R^2 = 0.47$; Table 4), followed closely by model residuals (partial $R^2 = 0.40$). For male sperm, transcriptional noise was positively correlated with treatment (P-value = 6.19×10^{-15} ; Fig. 5b), negatively correlated with gene expression (P-value=0; Fig. 5d) and gene body methylation (P-value = 8.45×10^{-50} ; Fig. 5f), and positively correlated with methylation variation (P-value = 6.40×10^{-163} ; Fig. 5h). Transcriptional noise was also negatively correlated with the interaction between treatment and gene expression and gene length, but positively correlated with gene expression squared, the interaction between treatment and gene expression squared, and the interaction between treatment and gene body methylation (Table 4). The model residuals had the highest partial R^2 (partial $R^2 = 0.62$) followed by methylation (partial $R^2 = 0.16$; Table 4). In both female gonads and male sperm, the statistically significant influence of pCO₂ treatment on transcriptional noise was associated with a small effect size, likely due to inter-sample variability.

Influence of methylation on alternative splicing

The influence of methylation was examined in the top 20 genes from PCs 5 and 6, which exhibited changes in alternative splicing from elevated pCO_2 exposure. There were no DMLs found in top female gonad genes, while six male sperm genes contained DMLs. Average gene body methylation in control and exposed samples was investigated as predictors of alternative splicing using sexspecific binomial GLMs. In both sexes, average gene body methylation in control samples, methylation in exposed samples, and gene length had no significant impact on alternative splicing status (Table 5; Fig. 6).

Discussion

As elevated pCO₂ continues to threaten marine invertebrates, it is critical to understand the role epigenetic and transcriptomic mechanisms play in stress resilience and the potential for environmental memory. Impacts on adult reproduction can influence intergenerational plasticity, so disentangling how various molecular mechanisms influence reproductive tissue is key for understanding offspring responses to stress. For example, herbicide exposure [34], sea surface temperature [35], OA [36], and OA and warming [37] have elicited methylation patterns that have subsequently been inherited by offspring. McNally et al. [27] used adult oysters to produce offspring and understand intergenerational plasticity in elevated pCO₂ conditions. After 3 days of exposure, larvae from parents exposed to elevated pCO₂ had higher shell growth rates and larger shell sizes, and these effects were pronounced when larvae were also reared in elevated pCO₂ conditions [27]. Interestingly, there was no impact of elevated pCO₂ on egg size or shape [27]. Since maternal provisioning may not have been responsible for larval phenotypes, exploring methylation and gene expression responses in gonadal tissues of those same adult oysters can elucidate regulatory mechanisms underlying observed carryover effects. By better understanding these mechanisms, we can better predict how environmental stress will influence intergenerational responses, and therefore ecosystem outcomes.

Given that the functional role of DNA methylation in marine invertebrates remains unclear [33], we leveraged whole genome, transcript-level data to conduct the first extensive assessment of

Table 4. Summary of multiple linear regression examining the correlation between the explanatory variables of gene expression, gene body methylation, the CV of gene body methylation, gene length, and pCO_2 treatment on the response variable of transcriptional noise, or the CV of gene expression

	Females				Males			
Predictor	eta Estimate	t-value	P-value	Partial R ²	β Estimate	t-value	P-value	Partial R ²
Intercept	0.14	19.71	3.04×10 ^{-86*}	_	0.43	19.22	1.22×10 ^{-81*}	_
Treatment	-8.62×10^{-3}	-3.17	0.002	0.02	0.15	7.81	6.19×10 ^{-15*}	3.96×10^{-3}
log ₂ (Gene expression) ²	0.01	35.85	1.10×10^{-278}	0.02	0.07	46.84	0	2.98×10^{-3}
log ₂ (Gene expression)	-0.10	-51.19	0*	0.06	-0.41	-47.87	0*	0.15
Gene methylation	-3.85×10^{-3}	-56.32	0	0.47	-1.40×10^{-3}	-14.88	$8.45 \times 10^{-50^*}$	0.16
log ₁₀ (CV methylation)	0.10	35.44	1.87 × 10 ^{-272*}	0.02	0.14	27.46	6.40×10 ^{-163*}	0.05
Treatment $\times \log_2(\text{Gene Expression})^2$	6.18×10^{-3}	14.89	4.81×10 ^{-50*}	2.80×10^{-3}	0.02	7.25	4.40×10^{-13}	2.29×10^{-6}
Treatment × log ₂ (Gene expression)	-0.07	-26.15	6.46×10^{-150}	4.29×10^{-3}	-0.10	-7.50	6.73×10 ^{-14*}	1.77×10^{03}
Treatment × Gene methylation	6.38×10^{-4}	7.43	1.13×10 ^{-13*}	3.06×10^{-3}	7.03×10^{-4}	7.06	1.67×10^{-12}	1.32×10^{-3}
log ₁₀ (Gene length)	7.14×10^{-3}	3.80	0.001*	9.90×10^{-5}	-0.03	-6.38	$1.87 \times 10^{-10^*}$	1.14×10^{-3}

Separate models were constructed for female gonads (adjusted R²=0.60) and male sperm (adjusted R²=0.38). Coefficients of determination, or partial R², were calculated separately for each predictor. *P-value<.05.



Figure 5. Relationship between transcriptional noise and (a, b) elevated pCO_2 treatment; (c, d) FPKM; (e, f) gene methylation (%); and (g, h) CV of methylation as defined by the best-fit linear model for female gonads (adjusted $R^2 = 0.60$) and male sperm (adjusted $R^2 = 0.38$) oysters.

DNA methylation–gene activity relationships in response to elevated pCO_2 in *Crassostrea* spp. gonad tissues to our knowledge. We found that DNA methylation was significantly correlated with gene expression variability, or transcriptional noise, in a sexspecific manner in response to elevated pCO_2 . These findings align with the growing consensus that gene body methylation may not have a direct regulatory role in gene-level expression, but instead functions to broadly reduce transcriptional variation across the genome [33, 36, 40, 55, 56]. Additionally, differences in gene activity and methylation patterns between female and male oysters highlight the need to examine how OA may affect reproduction differently in females and males [21] and underscore the cell- and tissue-specificity of DNA methylation [57, 58].

Methylation does not impact gene-level responses in gonads to elevated pCO₂

In this study, there was no substantial impact of elevated pCO_2 on gene-level expression in female gonads and male sperm. This is not surprising when considering a similar study on elevated pCO_2 on gene expression in *C. virginica* mantle tissue [42]. Additionally, the heterogeneity of cell types (eggs and connective tissue) in female samples could have increased expression variability and

reduced the ability to detect DEGs. There are several OA studies that do see an impact on gene expression [20], with the predominant functions of influenced genes associated with stress response, acid-base regulation, metabolic processes, and calcification. While we did not see substantial gene-level impacts, we did see significant changes to transcript-level activity—changes in the maximum number of transcripts expressed per gene, shifts in the predominant transcript expressed, and patterns of alternative splicing—and to DNA methylation. These changes were associated with biological processes such as reproduction (e.g. gamete development, regulation of female receptivity, and sperm motility), various signaling pathways, and protein ubiquitination. These data suggest the importance of these molecular mechanisms in oyster response to elevated pCO₂ conditions.

Beyond understanding the response of oyster reproductive tissue to elevated pCO_2 , a primary impetus for this work was to address hypotheses surrounding the role of methylation in environmental response and gene expression regulation. Following mammalian studies, it was previously considered that environmentally sensitive changes in methylation drive differential expressed genes on a gene-by-gene basis in marine invertebrates [59, 60]. However, to date, numerous molluscan studies have

Table 5. Binomial GLM results evaluating the influence of gene body methylation and gene length on alternative splicing status for the top 20 genes from each ASCA PC

	Females			Males		
Predictor	eta Estimate	z-value	P-value	eta Estimate	z-value	P-value
Intercept	4.80	1.67	0.09	-2.91	-1.25	0.21
Gene body methylation in control samples	-0.02	-0.44	0.65	-0.04	-1.03	0.30
Gene body methylation in control samples	-0.02	-0.37	0.71	0.04	1.14	0.25
log ₁₀ (Gene length)	-0.85	-1.26	0.2	0.49	0.85	0.39

No explanatory variables were significant predictors of alternative splicing status.



Figure 6. Average gene body methylation in control and exposed samples for genes that were alternatively spliced (difference) and not alternatively spliced (no difference) based on elevated pCO_2 exposure in (a) females and (b) males.

failed to find any significant relationships between differential DNA methylation and differential gene expression [42, 45, 49] or have found minimal overlaps in DEGs and differentially methylated regions [48, 61]. As described earlier, we could not examine the association between differential methylation and differential gene expression since we did not observe any of the latter. We also did not see a concordant change in gene expression at the gene level where differential methylation occurred. Taken together, our data suggest that changes in methylation within a gene do not drive concurrent changes in average gene expression in gonad

tissue, which is similar to what has been previously reported in molluscan somatic tissue [42, 45, 49].

Methylation regulates gene activity differently in female and male reproductive tissue

A critical regulatory role for DNA methylation is the reduction of gene expression variability, or transcriptional noise. We speculate that this genome-wide role may be more important than direct modulation of gene expression. This study found an inverse relationship between methylation and transcriptional noise: genes with higher levels of methylation had lower expression variation (Fig. 5). The relationship found in this study is consistent with previous research in molluscan somatic tissues [42, 45, 49, 62] and across invertebrate taxa [33, 55, 56]. We also demonstrate that methylation variability is positively correlated with transcriptional noise (Fig. 5). In other words, larger changes in gene body methylation are associated with higher expression variability. The general relationships described earlier were also consistent in response to elevated pCO₂, further supporting the idea that methylation has a broad regulatory role in response to stress. In this study, we show that these responses are also seen in gonad tissues. We also identified a sex-specific pattern: elevated pCO₂ decreases transcriptional noise in female gonads but increases noise in male sperm with respect to control conditions (Fig. 5). The difference in methylation-noise relationships between sexes could suggest that sperm cells require additional methylation to achieve similar levels of transcriptional noise reductions to females. As egg quality is essential to successful reproduction, reducing transcriptional noise may be necessary to maintain egg quality, while gene expression variability may not be maladaptive for sperm. Additionally, eggs contain higher levels of RNA than sperm [63], so modulating transcriptional noise may be more important in eggs. Given the small effect size of pCO₂ treatment on transcriptional noise in males, future work should investigate the phenotypic implications of these shifts. Combined with the subtle gene expression differences between control and OA-exposed oysters, our work suggests that reductions in general gene variability are an integral component of OA response in gonad tissues of C. virginica.

Additional differences in female and male reproductive tissue responses to elevated pCO₂ underscore the importance of disentangling biological and environmental effects on gene expression and DNA methylation. It is well known that OA can have sex-specific impacts on physiology [21], and previous work in C. gigas has documented distinct gene expression patterns associated with oogenesis and spermatogenesis [53]. We extend this work by demonstrating contrasting OA effects between female gonads and male sperm. We found that sperm had more genes with higher number of transcripts expressed in elevated pCO₂, as opposed to female gonad tissue that had more genes with a smaller number of transcripts expressed in the same conditions (Table 1). Even though OA-responsive patterns of alternative splicing were similar between sexes, the top 20 gene drivers of those patterns differed and were involved in sex-specific reproductive processes. For example, protein UFMylation genes were alternatively spliced in female gonad tissue, while genes involved in regulation of metallopeptidase activity were alternatively spliced in male sperm. Protein UFMylation is involved in broader protein ubiquitination processes, which are conserved across oyster responses to pCO₂ [11, 43] and characteristic of female gonad methylome responses [46]. Metallopeptidases are important for sperm motility in humans [64, 65] and have been found to increase in expression in C. gigas during spermatogenesis [66].

Differences in female and male reproductive tissue responses extend to methylation as well. Sperm had a more dramatic methylation response to elevated pCO_2 , with 2916 DMLs compared to 89 in female gonad tissue oysters, even though a similar number of CpGs with 10× coverage were detected in both sexes. Sexspecific differences in methylation [50-52] are well documented in C. gigas. Male C. gigas have higher levels of gene body methylation than females in both somatic [52] and reproductive tissue [50, 51]. Our results confirm that these patterns are present in C. virginica gonads as well. These gene expression and methylation differences may be attributed to cell type: oocytes typically have higher levels of maternal RNA as opposed to sperm cells with more DNA [63]. Enrichment of processes such as oocyte development in females and spermatogenesis in males in genes with altered activity or with DMLs align with the specific biological roles of these cell types. In our study, female samples were likely a mix of oocytes and connective tissue, which further complicates the observed patterns. Previous studies have documented environmentally sensitive methylation in response to elevated pCO₂ in molluscan reproductive tissue [44, 46], but none examine sex-specific patterns or have data for both sexes. As the role of methylation is cell-, tissue-, and sex-specific [57, 58], future investigations should compare molecular and phenotypic responses to elevated pCO₂ between tissues and sexes when appropriate.

Methylation alters some aspects of transcript-level gene activity in gonad tissues

Although there is limited evidence for DNA methylation driving differential gene expression in molluscs, it is possible that methylation may change transcriptional opportunities. Sparse methylation may allow for exon skipping or access to alternative start sites, thereby modifying gene expression and phenotype [32, 33, 67]. Using differential exon expression as a proxy for alternative splicing, we observed that elevated pCO₂ changes exon expression patterns (Figs 1 and 2). However, environmentally responsive splicing accounted for a small proportion for the total splicing variance. There were minimal overlaps with DMLs and alternatively spliced genes, and gene body methylation did not significantly influence splicing (Table 5). This contrasts with work in C. gigas and upwelling-exposed Strongylocentrotus purpuratus urchin larvae, showing that high levels of methylation in exons [68, 69] and/or introns [69] are associated with alternative splicing. In C. gigas, alternative splicing was explored in one gene (CGI_10021620) with higher methylation and gene expression in mantle tissues as opposed to male gametes [68]. The authors conclude that lower levels of methylation in male gametes may have opened access to an alternative stop codon, producing a shorter protein product when compared to the product from mantle tissues [68]. Since C. gigas were not exposed to any stressors in this previous study, the role of methylation in regulating alternative splicing could be distinct for cell-type differentiation versus environmental response. Significant methylation regulation of alternative splicing was found in S. purpuratus larvae from female urchins exposed to upwelling, but only when considering transcriptional start site accessibility [69]. It is possible that incorporating additional epigenetic mechanisms like chromatin state, or utilizing long-read sequencing methods for gene expression data, would elucidate connections between methylation and alternative splicing in C. virginica.

The lack of correlation between methylation and alternative splicing in our study could indicate that these molecular mechanisms are not intrinsically linked in *C. virginica*. In both sexes,

we observed distinct biological functions associated with alternatively spliced genes and genes containing DMLs. For example, in female oysters alternatively spliced genes were involved in metabolic and developmental processes such as the glycolysis pathway and DNA ligation, while genes with DMLs had prominent roles in apoptosis, reproductive processes, and cell organization and biogenesis. The latter processes have been previously implicated in oyster response to elevated pCO₂ [11, 43], including studies examining gonad methylome responses [44, 46]. Another potential explanation is that methylation is decoupled from transcript activity in gametes. The majority of RNA in oocytes is acquired during vitellogenesis, which may not reflect gene expression responses to OA [63]. On the other hand, sperm cells have lower RNA content and are not as transcriptionally active [63]. Therefore, it is possible that DNA methylation and alternative splicing have separate, but essential, roles in moderating reproductive tissue response to OA.

Although methylation and alternative splicing patterns were not correlated, the influence of gene body methylation on other aspects of gene activity suggests a potential regulatory role. In male oysters, methylation was significantly associated with changes in the maximum number of transcripts expressed per gene and was marginally associated with shifts in the predominant transcripts (Tables 2 and 3). Interestingly, genes both containing DMLs and exhibiting alterations to gene activity had similar enriched biological processes involved in signal transduction and cell organization and biogenesis. These commonalities further support the idea methylation may have a role in gene activity beyond regulating alternative splicing patterns for specific pathways.

Conclusion

Elucidating the relationship between epigenetic and transcriptomic mechanisms will allow for a holistic understanding of phenotypic responses to climate stressors. We highlight the importance of transcript-level data for parsing the relationship between these mechanisms. Future research should integrate transcriptlevel data along with exon- and intron-specific methylation and expression analyses to further clarify the relationship between methylation and gene activity within and across generations. As DNA methylation is a potential mechanism for intergenerational carryover effects [34–37], it is critical to assess environmentally responsive methylation in adult reproductive tissue. Our study suggests that methylation has a genome-wide regulatory role, effectively maintaining gene expression homeostasis in reproductive tissues under elevated pCO₂ by reducing transcriptional noise. Importantly, we found that the relationship between methylation and transcriptional noise was different between female reproductive tissue and sperm: male oysters required higher levels of methylation to achieve similar reductions in transcriptional noise when compared to female oysters. An accompanying study using the same adult C. virginica found no difference in female egg quality after OA exposure during reproductive conditioning, and no delays in early larval development [27]. Since this signal was observed even with significant genotype effects [27], it suggests that epigenetic maintenance of reproduction may also confer intergenerational resilience to elevated pCO_2 . We demonstrate that OA conditions lead to molecular changes in oyster gonads, and future work should evaluate specific connections between these changes and offspring fitness to understand whether inheritance of environmental memory can occur.

Materials and methods Experimental design and seawater chemistry analysis

The experimental design and seawater chemistry manipulations have been published by McNally *et al.* [27]. Briefly, adult *C. virginica* (mean \pm SD shell length: 7.92 \pm 1.80 cm) were acclimated to ambient pCO₂ conditions (mean pCO₂ \pm SD = 632 \pm 64 ppm) for 1 week prior to random placement in experimental pCO₂ conditions [either control (572 \pm 107 ppm) or elevated pCO₂ (2827 \pm 360 ppm)] during reproductive conditioning at 20°C. The elevated pCO₂ treatment corresponds with seawater conditions undersaturated with aragonite and is consistent with observations in estuarine ecosystems that oysters inhabit [42, 70]. Additionally, the elevated pCO₂ treatment was chosen to increase the inferential and statistical power of the study [71].

Four replicate 42-l tanks containing 10 oysters each were used for each pCO₂ condition. Oysters were exposed to experimental conditions for 30 days, which has shown to elicit differences in C. virginica gonad methylation [44]. A flow-through OA array was used to control temperature, salinity, and pCO2 with temperature, salinity, and pH_T recorded three times per week during the experimental period. Seawater samples were collected during the first and third weeks of the experiment for dissolved inorganic carbon and total alkalinity measurements. Oysters were fed in accordance with best practices from Helm and Bourne [72]. No adult mortalities or significant random effects of adult tank on egg quality were reported [27]. At the end of the 30-day exposure, gonad samples from adult oysters were visually inspected under a microscope to determine sex and maturity, with sperm samples also checked for motility, targeting at least eight mature females and eight mature males per treatment [27]. Gonad tissues were collected from 16 female oysters (eight from three replicate control tanks and eight from two replicate elevated pCO₂ tanks), and motile sperm from 10 male oysters (four control and six exposed to elevated pCO_2), with at least one individual from each replicate tank, for WGBS and RNA-seq.

Nucleic acid extraction and library preparation

Simultaneous DNA and RNA extraction was conducted for female gonad and sperm samples using the Zymo Quick-DNA/RNA Microprep Plus Kit (Cat # D7005). A total of 10–15 mg of frozen gonad tissue and 50 μ l of sperm in seawater solution were used as inputs for females and males, respectively. The manufacturer's instructions were used with the following exceptions. After addition of 300 μ l DNA/RNA Shield (1×), 30 μ l PK Digestion Buffer, and 15 μ l Proteinase K, female samples were incubated at 55°C overnight. Four volumes of DNA/RNA lysis buffer (200 μ l) were added to sperm samples. All centrifugation steps were conducted at 16 000 rcf. DNase I digestion was performed for RNA. Concentrations of eluted DNA and RNA were verified with Qubit dsDNA high sensitivity or RNA high sensitivity assays (Thermo Fisher Scientific), respectively, and quality confirmed on a BioAnalyzer.

All samples were submitted to ZymoResearch for library preparation and sequencing. WGBS libraries were created from 100 ng of DNA with the Zymo-Seq WGBS Library Kit (Cat # D5465) following the manufacturer's protocol and PCR performed with Illumina Unique Dual Indices. Library quality was checked with an Agilent 2200 TapeStation. Libraries were sequenced on an Illumina NovaSeq instrument to generate 150-bp paired-end reads. Prior to sequencing, total RNA-seq libraries were created using 250 ng of RNA. For all samples, rRNA was removed following methods described in Bogdanova *et al.* [73] with some modifications. Libraries were constructed using the Zymo-Seq RiboFree Total RNA Library Prep Kit (Cat # R3000) according to the manufacturer's instructions. Libraries were sequenced on an Illumina NovaSeq to generate at least 30 million read pairs for 150-bp paired-end sequences per sample. Resulting sequences were submitted to the National Center for Biotechnology Information (NCBI) Sequencing Read Archive under BioProject accession number PRJNA1089235.

Genome information and feature tracks

The C. virginica genome was used for several analyses [74]. Gene, CDS, exon, and lncRNA tracks were provided by the C. virginica genome NCBI RefSeq annotation (RefSeq: GCF_002022765.2) on NCBI [74]. These tracks were used to generate the remaining feature tracks with BEDtools v2.26.0 [75]. The complement of the exon track, or noncoding sequences, was created with BEDtools complementBed v.2.26.0. Overlaps between these noncoding sequences and genes-introns-were obtained using intersectBed. Exon UTR were identified by subtracting the coding sequence track from the exon track with subtractBed. Upstream flanking regions, or putative promoters, were defined as the 1000 bp upstream of a transcription start site. These flanks were created by adding 1000 bp upstream of genes with flankBed, taking strandedness into account. Other genes that overlapped with these flanks were removed using subtractBed. Flanks 1000 bp downstream of genes were created as well. To identify intergenic regions, the complement of the gene track was created with complementBed, then subtractBed was used to remove flanking regions from the intergenic region track. TE information from RepeatMasker was obtained from the NCBI RefSeq annotation [76, 77]. CG motifs in the C. virginica genome were identified using the fuzznuc function from EMBOSS. The number of CG motifs in a given track were obtained using intersectBed between the CG motifs and a feature track. The genome features are available along with corresponding code [78].

Genetic variation

SNPs were identified in the BS-seq data using the Nextflow [79] EpiDiverse/SNP pipeline [80] to characterize genetic differences between samples. Sorted, deduplicated BAM files, generated through methylation analysis with Bismark [81], were used as inputs, along with the NCBI *C. virginica* genome FastA (GCF_002022765.2_C_virginica-3.0_genomic.fa) provided as the reference. The options –variants and –clusters were provided when running the EpiDiverse/SNP pipeline with default settings (–min-coverage 0, –min-repeat-entropy 1). After variant identification, NgsRelate was used to estimate pairwise relatedness using Variant Call Format file output from EpiDiverse/SNP [82]. Variants were filtered to include only bi-allelic sites (–min-alleles 2 –max-alleles 2) and minor allele counts of at least two (–mac 2). Variants were included if data were missing for >50% of the samples (–max.missing 0.5).

Recent studies have shown that genetic population structure can influence methylation [83]. In order to determine if genetic variation had a significant impact on gene expression and methylation, we first performed correlation analyses to determine whether genetic distance correlated with methylation and gene expression distance between individuals. In the case of a significant correlation, genetic distance would need to be included in downstream model analyses of differential methylation and gene expression. To perform these correlations, we generated Euclidean distance matrices of genetic data, expression data, and DNA methylation data. For genetic data, SNPs were derived from the Nextflow EpiDiverse SNP Pipeline as described earlier. Gene expression matrices were derived from DESeq2 output [84]. DNA methylation data were obtained from methylkit output using a 10× coverage minimum threshold. For analysis, Mantel tests (mantel.rtest) in R were used with 99 replicates. We visualized these relationships using Pearson's correlation matrices. Corresponding code is available for all genetic variation analyses [78].

Gene activity

Gene and transcript expression

FastQ files were quality trimmed using fastp [85] default settings for paired-end adapter trimming, followed by the removal of the last 20 bases from the five prime ends of all reads. These bases were removed because the FastQC "Per Base Sequence Content" assessment showed these exhibited a less uniform distribution of nucleotides than the remainder of the read length. Trimmed reads were then visualized with MultiQC [86]. To identify isoforms, the C. virginica genome (NCBI accession: GCF 002022765.2) was indexed using HISAT2 [87]. A pipeline detailed by Pertea et al. [88] was used as a guide to identify isoforms and analyze differential expression. Trimmed reads were aligned to the genome using HISAT2 (v2.1.0), followed by StringTie [89] to identify isoforms and generate files for downstream analysis using the R [90] packages ballgown [91] and tidyverse [92]. DEGs and DETs (P-value and q-value < .05) were identified between pCO₂ treatments within a given sex using ballgown [91], with "FPKM" as the expression measurement.

Elevated pCO_2 may influence finer-scale expression processes. To explore this, we quantified differences in the maximum number of transcripts expressed in each gene for each sex separately. The maximum number of transcripts for each gene was identified across all samples. The difference in the maximum number of transcripts expressed in each gene was calculated with respect to the elevated pCO_2 -exposed samples. Positive differences signify more unique transcripts expressed in elevated pCO_2 conditions, while the opposite holds true for negative differences.

An enrichment test with topGO [93] was conducted to understand if certain biological processes were over-represented in genes with differences in the maximum number of transcripts expressed. First, a gene ID-to-GO term database (geneID2GO) was created for manual GO term annotation. Each line of the database contained gene ID in one column, and all corresponding GO terms in another. The list of gene IDs from this database was used as the gene universe for enrichment. Two separate lists were compiled for genes of interest: one for genes with increased transcript counts and another for genes with decreased transcript counts due to elevated pCO₂ exposure. A topGO object was generated for each sex- and direction-specific list of genes and biological process GO terms, with GO term annotation performed using the geneID2GO database. A Fisher's exact test was used to identify GO terms significantly enriched with respect to the gene background (P-value < .01). Corresponding code is available for all gene activity analyses [78].

Predominant transcript identification

In addition to quantifying transcript expression, we examined shifts in the predominant transcript—the transcript with the highest mean level of expression in an experimental group—due to low pH exposure within male and female samples separately. Gene expression data from ballgown were filtered to retain genes with multiple transcripts where gene expression was >0. Differences in the identity of the predominant transcript were determined for genes remaining after filtering. An enrichment test with topGO [93] was used to identify significantly enriched biological process GO terms (P-value < .01) in genes where the predominant transcript changed using similar methods as described earlier (see "Gene and transcript expression").

Alternative splicing

The final way we characterized gene activity was through an alternative splicing analysis. We hypothesized that OA exposure would induce different patterns of exon expression, or alternative splicing. To characterize these patterns, the expression of exons 2-6 was calculated as fold change values relative to the expression of exon 1 for every gene in each sample. To do this, bedtools::coverage was used with BAM alignment files from HISAT2 and the genome feature file. A minimum read count coverage of 10× was used. These values were generated for females and males separately. We then used ASCA using the ALASCA package (v1.0.14) in R [94] to perform longitudinal multivariate analysis across exon positions. ASCA is useful for the analysis of both fixed and random effects on high-dimensional multivariate data when other multivariate ANOVA analyses (i.e. MANOVA) are not able to include more variables than there are observations [54]. Each sex-specific ASCA model analyzed multivariate expression as a function of exon position, treatment, and their interaction, with the sample as a random effect to account for repeated measures:

```
expression relative to exon 1 \sim exon position * treatment + (1|sample)
```

Genes that were detected in all samples were included (11270 genes in females and 12645 genes in males). The model was validated with bootstrapping with replacement for 100 iterations [95]. We extracted and plotted PCs that explained >1% of the variance to visualize expression patterns across exons. We categorized PCs as those that either (I) show no difference in exon expression patterns between treatments or (II) show differences between treatments. For each PC, we extracted the top 20 genes with the highest PC scores. These gene lists were combined into the two previously described categories for downstream analyses. For these top genes, the number of genes with changes in the maximum number of transcripts expressed were quantified for females and males.

A functional enrichment analysis was conducted with topGO [93] to understand if alternatively spliced genes contained any over-represented biological processes. Enrichment was conducted using similar methods as described earlier (see "Gene and transcript expression"). All genes included in the ASCA were used as the gene background for this analysis. Biological process GO terms with P-values < .01 were considered significant.

DNA methylation analysis CpG locus-level methylation quantification

Quality-trimmed sequence reads were aligned to the C. virginica genome using Bismark v.0.22.3 with -score_min L,0,-0.6 to dictate alignment specificity. In order to characterize CpG methylation for individual samples, bedGraph files were generated based on Bismark coverage2cytosine output, where CpG data from each strand were merged (-merge_CpG), filtered to retain CpG loci with at least 10× coverage, and sorted. Since $C \rightarrow T$ SNPs can be interpreted as unmethylated cytosines, they were identified in bedGraphs using default BS-SNPer settings [96], then the methylation values of those $C \rightarrow T$ SNPs were changed to 0.

Sex-specific baseline methylation landscapes were characterized using methylation information for CpG loci with at least $10 \times$ coverage in a sample. Individual bedGraph files (described earlier) were combined using bedtools::unionBedGraphs v2.30.0 [75] to create sex-specific union bedGraphs. Percent methylation was averaged across all samples for each CpG locus. Highly (>50%), moderately (10–50%), and lowly (\leq 10%) methylated CpGs were identified in sex-specific union bedGraphs. Overlaps between highly methylated CpGs and various genome features were characterized using bedtools::intersectBed. A chi-squared contingency test (chi.sq) was used to determine if the distribution of highly methylated CpGs was significantly different from the distribution of all CpGs in the *C. virginica* genome, with P-values < .05 indicating a significant difference.

Deduplicated BAM alignment files from Bismark were used for sex-specific differential methylation analysis with methylKit [97]. The threshold for comparison was a minimum of 10× coverage (lo.count = 10) and a maximum coverage set at 98% (hi.perc = 98), using the destranded setting (destrand = TRUE). Putative DMLs were determined based on a 50% difference in the comparison group and a q-value threshold of .01. Finally, $C \rightarrow T$ SNPs were removed from putative female and male DML lists using bedtools::subtractBed. Genomic locations of female and male DMLs were determined by overlapping with annotated features using bedtools::intersectBed. A chi-squared contingency test was used to test the null hypothesis of no association between methylation and specific genome features using all CpGs with 10× data and DMLs for females and males separately. A P-value of <.05 signified a significantly different proportion of DMLs in a specific genome feature when compared to 10× CpG data. An enrichment test with topGO [93] was used to identify significantly enriched biological process GO terms (P-value < .01) in genes containing DMLs. The gene universe for the enrichment was a sex-specific list of genes containing 10× CpGs. The enrichment was conducted using similar methods as described earlier (see "Gene and transcript expression").

Gene-level methylation quantification

Gene-level methylation was determined by taking the mean value of methylation at loci within a gene where coverage was at least 10×. Specifically, bedGraph files generated as part of the Bismark alignment described earlier were used in conjunction with bedtools::intersectBed to extract CpG loci for each gene in each sample.

Methylation influence on gene activity

To broadly examine methylation and gene expression signatures of OA treatment for each sex, we used a similar approach to the genetic variation analysis. We used Monte Carlo tests (mantel.rtest) to calculate correlations between Bray–Curtis distance matrices derived from gene methylation and expression datasets. Separate subsequent analyses were conducted to examine if methylation influenced specific aspects of gene activity.

Influence of methylation on gene and transcript expression

The first set of analyses was used to determine the relationship between gene methylation and expression. First, the number of DMLs in DEGs and DETs were quantified. Next, we investigated if gene methylation influenced differences in the maximum number of transcripts expressed between control and exposed samples using a binomial GLM (glm). For each sex separately, we examined if a change in the maximum number of transcripts expressed due to OA exposure ("success") could be explained by change in gene body methylation, number of DML in a gene, change in gene expression, or gene length: occurrence of a change in the maximum number of expressed transcripts \sim change in gene methylation + number of DML in the gene+log2(change in gene

Change in gene body methylation was calculated by subtracting average gene body methylation in the control treatment from the OA treatment. A similar calculation was done to obtain change in gene expression. Explanatory variables were considered significant if P-values < .05. Corresponding code is available for all gene activity analyses [78].

Influence of methylation on predominant transcript shifts

The second set of analyses examined if previously identified transcript shifts were due in part to gene methylation. Similar sex-specific binomial GLM (glm) to those described earlier (see "Influence of methylation on gene and transcript expression") was constructed to examine if the occurrence of a predominant transcript shift due to OA exposure could be explained by change in gene body methylation, number of DML in a gene, change in gene expression, or gene length:

occurrence of predominant transcript shift \sim change in gene methylation+number of DML in the gene+log2 (change in gene expression+1)+log10(gene length)

Predictors were significant if P-values < .05.

Transcriptional noise analysis

Following Wu *et al.* [98], transcriptional noise, or the CV of gene expression, was used as a response variable in a linear model (lm) with average gene methylation, average gene expression, CV of gene methylation, gene length, OA treatment, and the interaction between treatment and methylation or expression:

 $\label{eq:constraint} \begin{array}{l} \log \$_10\$ (\text{CV of gene expression}) \sim \text{gene methylation} + \\ \log \$_2\$ (\text{gene expression}+1)\$^2\$ + \log \$_2\$ (\text{gene} \\ \text{expression}) + \log \$_10\$ (\text{CV of gene methylation}) + \\ \log \$_10\$ (\text{gene length}) + \text{treatment} + \text{treatment} \ast \text{gene} \\ \text{methylation} + \text{treatment} \ast \log \$_2\$ (\text{gene expression}+1) \\ \$^2\$ + \text{treatment} \ast \log \$_2\$ (\text{gene expression}+1) \\ \end{array}$

The polynomial terms for gene expression were included after initial data visualization and represent a relationship between gene expression and transcriptional noise where gene expression variability is the highest at very low or high levels of gene expression. A backward deletion approach was used for model construction, with explanatory variables removed from the model based on a *P*-value of <.05 threshold. Competing models were evaluated with the Akaike information criterion (AIC). The best model fit was defined by the lowest AIC and all input variables having a *P*-value of <.05. Residuals of the best-fit model were evaluated for normality and homoscedasticity. Coefficients of determination, or partial R^2 , were calculated for all significant explanatory variables. To calculate partial R^2 , the sum of squares for all input variables and residuals.

Influence of methylation on alternative splicing

Finally, methylation's influence on gene activity was evaluated by understanding methylation patterns in alternatively spliced genes. The number of DMLs were counted in genes that were and were not alternatively spliced. A sex-specific binomial GLM (glm) was used to determine if average gene body methylation in control whether or not the gene was alternatively spliced \sim average gene body methylation in control samples + average gene body methylation in exposed samples + log10(gene length)

Terms were considered significant predictors of alternative splicing if P-values <.05.

Acknowledgements

We are grateful to Grace Crandall for assistance with DNA and RNA extractions and to Drs Shayle Matsuda, Katherine Silliman, Laura H. Spencer, and Samuel N. Bogan for insight on manuscript ideation and earlier drafts of this manuscript.

Author contributions

A.D.-W. and K.E.L. conceived and ran the experiment. Y.R.V. performed DNA and RNA extractions. All authors contributed to initial analyses. Y.R.V., A.S.H., S.J.W., and S.B.R. conducted final analyses. A.D.-W. and K.E.L. contributed to analysis. Y.R.V., A.S.H., S.J.W., and S.B.R. wrote the initial manuscript draft. All authors reviewed and approved the manuscript.

Supplementary data

Supplementary data is available at EnvEpig online.

Conflict of interest: None declared.

Funding

This work was facilitated through the use of advanced computational, storage, and networking infrastructure provided by the Hyak supercomputer system at the University of Washington. This work was funded by the National Science Foundation (award no. 1634167 to S.B.R. and award no. 1635423 to K.E.L.), by a grant from Washington Sea Grant, University of Washington, pursuant to National Oceanic and Atmospheric Administration award no. NA22OAR4170103, and by the United States Department of Agriculture National Institute of Food and Agriculture National Animal Genome Research Program (NRSP-8) Small Funding Opportunity in Aquaculture to SAW and SBR. The views expressed herein are those of the authors and do not necessarily reflect the views of funding agencies.

Data Availability

Raw WGBS and RNA-seq data can be accessed at the NCBI Sequence Read Archive under BioProject accession number PRJNA1089235. Additional data, genome feature tracks, scripts, and supplementary materials are available in the *C. virginica* methylation and gene expression repository on the Open Science Framework (https://osf.io/xuy2f/) and on GitHub (https://github. com/sr320/ceabigr) [78].

References

1. Calvin K, Dasgupta D, Krinner G *et al.* IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. Arias P, Bustamante M, Elgizouli I *et al.* (eds.). Intergovernmental Panel on Climate Change (IPCC), 2023.

- 2. Gazeau F, Parker LM, Comeau S et al. Impacts of ocean acidification on marine shelled molluscs. *Mar* Biol 2013;**160**:2207–45.
- Ducker J, Falkenberg LJ. How the Pacific oyster responds to ocean acidification: development and application of a meta-analysis based adverse outcome pathway. Front Mar Sci 2020;7:597441.
- Parker LM, Ross PM, O'Connor WA et al. Predicting the response of molluscs to the impact of ocean acidification. Biology 2013;2:651–92.
- Lemasson AJ, Fletcher S, Hall-Spencer JM et al. Linking the biological impacts of ocean acidification on oysters to changes in ecosystem services: a review. J Exp Mar Biol Ecol 2017;492:49–62.
- Gazeau F, Quiblier C, Jansen JM et al. Impact of elevated CO₂ on shellfish calcification. Geophys Res Lett 2007;34:L07603.
- 7. Kurihara H, Kato S, Ishimatsu A. Effects of increased seawater pCO_2 on early development of the oyster Crassostrea gigas. Aquat Biol 2007;**1**:91–98.
- 8. Beniash E, Ivanina A, Lieb NS *et al*. Elevated level of carbon dioxide affects metabolism and shell formation in oysters *Crassostrea virginica*. *Mar Ecol Prog Ser* 2010;**419**:95–108.
- Ries JB. A physicochemical framework for interpreting the biological calcification response to CO₂-induced ocean acidification. *Geochim Cosmochim Acta* 2011;**75**:4053–64.
- Tomanek L, Zuzow MJ, Ivanina AV et al. Proteomic response to elevated pCO₂ level in eastern oysters, Crassostrea virginica: evidence for oxidative stress. J Exp Biol 2011;214:1836–44.
- 11. Timmins-Schiffman E, Coffey WD, Hua W et al. Shotgun proteomics reveals physiological response to ocean acidification in *Crassostrea gigas. BMC Genom* 2014;**15**:951.
- 12. Dineshram R, Chandramouli K, Ko GWK et al. Quantitative analysis of oyster larval proteome provides new insights into the effects of multiple climate change stressors. *Global Chang Biol* 2016;**22**:2054–68.
- Boulais M, Chenevert KJ, Demey AT et al. Oyster reproduction is compromised by acidification experienced seasonally in coastal regions. Sci Rep 2017;7:13276.
- 14. Omoregie E, Mwatilifange NSI, Liswaniso G. Futuristic ocean acidification levels reduce growth and reproductive viability in the pacific oyster (*Crassostrea gigas*). J Appl Sci Environ Manage 2019;**23**:1747–54.
- 15. Coen LD, Brumbaugh RD, Bushek D et al. Ecosystem services related to oyster restoration. *Mar Ecol Prog Ser* 2007;**341**:303–07.
- Smith RS, Cheng SL, Castorani MCN. Meta-analysis of ecosystem services associated with oyster restoration. *Conserv Biol* 2023;**37**:e13966.
- 17. Grabowski JH, Brumbaugh RD, Conrad RF et al. Economic valuation of ecosystem services provided by oyster reefs. *Bioscience* 2012;**62**:900–09.
- Kroeker KJ, Kordas RL, Crim RN et al. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. Ecol Lett 2010;13:1419–34.
- Melzner F, Mark FC, Seibel BA et al. Ocean acidification and coastal marine invertebrates: tracking CO₂ effects from seawater to the cell. Ann Rev Mar Sci 2019;12:499–523.
- Strader ME, Wong JM, Hofmann GE. Ocean acidification promotes broad transcriptomic responses in marine metazoans: a literature survey. Front Zool 2020;17:7.
- 21. Padilla-Gamiño JL, Alma L, Spencer LH et al. Ocean acidification does not overlook sex: review of understudied effects and

implications of low pH on marine invertebrate sexual reproduction. Front Mar Sci 2022;**9**:977754.

- 22. Gienapp P, Teplitsky C, Alho JS et al. Climate change and evolution: disentangling environmental and genetic responses. Mol Ecol 2008;17:167–78.
- 23. Byrne M, Foo SA, Ross PM et al. Limitations of cross- and multigenerational plasticity for marine invertebrates faced with global climate change. Global Chang Biol 2020;26: 80–102.
- 24. Parker LM, Ross PM, O'Connor WA et al. Adult exposure influences offspring response to ocean acidification in oysters. *Global Chang* Biol 2012;**18**:82–92.
- 25. Parker LM, O'Connor WA, Raftos DA et al. Persistence of positive carryover effects in the oyster, *Saccostrea glomerata*, following transgenerational exposure to ocean acidification. PLoS One 2015;**10**:e0132276.
- 26. Spencer LH, Venkataraman YR, Crim R et al. Carryover effects of temperature and pCO₂ across multiple Olympia oyster populations. Ecol Appl 2019;**30**:e02060.
- 27. McNally EM, Downey-Wall AM, Titmuss FD et al. Parental exposure of Eastern oysters (*Crassostrea virginica*) to elevated pCO₂ mitigates its negative effects on early larval shell growth and morphology. *Limnol Oceanogr* 2022;**67**:1732–45.
- 28. Griffith AW, Gobler CJ. Transgenerational exposure of North Atlantic bivalves to ocean acidification renders offspring more vulnerable to low pH and additional stressors. Sci Rep 2017;7:11394.
- 29. Venkataraman YR, Spencer LH, Roberts SB. Larval response to parental low pH exposure in the pacific oyster *Crassostrea gigas*. J Shellfish Res 2019;**38**:743.
- Deans C, Maggert KA. What do you mean, "Epigenetic"? Genetics 2015;199:887–96.
- 31. Bird A. DNA methylation patterns and epigenetic memory. *Genes* Dev 2002;**16**:6–21.
- 32. Eirin-Lopez JM, Putnam HM. Marine environmental epigenetics. Ann Rev Mar Sci 2018;**11**:335–68.
- 33. Bogan SN, Yi SV, Hoffmann F. Potential role of DNA methylation as a driver of plastic responses to the environment across cells, organisms, and populations. *Genome Biol Evol* 2024;**16**:evae022.
- 34. Rondon R, Grunau C, Fallet M et al. Effects of a parental exposure to diuron on Pacific oyster spat methylome. Environ Epigenet 2017;**3**:dvx004.
- 35. Liew YJ, Howells EJ, Wang X et al. Intergenerational epigenetic inheritance in reef-building corals. Nat Clim Chang 2020;10:254–59.
- 36. Lee YH, Kim M-S, Wang M *et al.* Epigenetic plasticity enables copepods to cope with ocean acidification. Nat Clim Chang 2022;**12**:918–27.
- Brennan RS, deMayo JA, Finiguerra M et al. Epigenetic and evolutionary mechanisms uniquely contribute to rescue from global change. bioRxiv 2024:2024-03.
- Peterson CR, Scott CB, Ghaffari R et al. Mixed patterns of intergenerational DNA methylation inheritance in Acropora. Mol Biol Evol 2024;41:msae008.
- 39. Bogan SN, Johnson KM, Hofmann GE. Changes in genome-wide methylation and gene expression in response to future pCO₂ extremes in the antarctic pteropod Limacina helicina antarctica. Front Mar Sci 2020;6:788.
- 40. Liew YJ, Zoccola D, Li Y *et al.* Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building coral. *Sci Adv* 2018;**4**:eaar8028.
- 41. Strader ME, Kozal LC, Leach TS *et al*. Examining the role of DNA methylation in transcriptomic plasticity of early stage sea

urchins: developmental and maternal effects in a kelp forest herbivore. Front Mar Sci 2020;**7**:205.

- 42. Downey-Wall AM, Cameron LP, Ford BM et al. Ocean acidification induces subtle shifts in gene expression and DNA methylation in mantle tissue of the eastern oyster (*Crassostrea virginica*). Front Mar Sci 2020;**7**:828.
- **43**. Lim Y-K, Cheung K, Dang X *et al*. DNA methylation changes in response to ocean acidification at the time of larval metamorphosis in the edible oyster, *Crassostrea hongkongensis*. *Mar Environ Res* 2020;**163**:105214.
- 44. Venkataraman YR, Downey-Wall AM, Ries J et al. General DNA methylation patterns and environmentally-induced differential methylation in the eastern oyster (*Crassostrea virginica*). Front Mar Sci 2020;**7**:225.
- Chandra Rajan K, Yuan M, Yu Z et al. Oyster biomineralisation under ocean acidification: from genes to shell. Global Chang Biol 2021;27:3779–97.
- 46. Venkataraman YR, White SJ, Roberts SB. Differential DNA methylation in Pacific oyster reproductive tissue in response to ocean acidification. BMC Genom 2022;23:556.
- 47. Dang X, Lim Y-K, Li Y et al. Epigenetic-associated phenotypic plasticity of the ocean acidification-acclimated edible oyster in the mariculture environment. Mol Ecol 2023;32:412–27.
- 48. Johnson KM, Sirovy KA, Casas SM et al. Characterizing the epigenetic and transcriptomic responses to Perkinsus marinus infection in the eastern oyster Crassostrea virginica. Front Mar Sci 2020;7:598.
- 49. Johnson KM, Sirovy KA, Kelly MW. Differential DNA methylation across environments has no effect on gene expression in the eastern oyster. J Anim Ecol 2021;91:1135–47.
- 50. Sun D, Li Q, Yu H. DNA methylation differences between male and female gonads of the oyster reveal the role of epigenetics in sex determination. *Gene* 2022;**820**:146260.
- 51. Sun D, Yu H, Kong L et al. The role of DNA methylation reprogramming during sex determination and sex reversal in the Pacific oyster Crassostrea gigas. Int J Biol Macromol 2024;259:128964.
- 52. Teng W, Xu C, Liu S et al. DNA methylation in oysters is influenced by genetics and sex. Res Square 2024.
- 53. Broquard C, Saowaros S-A, Lepoittevin M et al. Gonadal transcriptomes associated with sex phenotypes provide potential male and female candidate genes of sex determination or early differentiation in Crassostrea gigas, a sequential hermaphrodite mollusc. BMC Genom 2021;22:609.
- 54. Bertinetto C, Engel J, Jansen J. ANOVA simultaneous component analysis: a tutorial review. Anal Chim Acta X 2020;**6**:100061.
- 55. Sarda S, Zeng J, Hunt BG *et al*. The evolution of invertebrate gene body methylation. Mol Biol Evol 2012;**29**:1907–16.
- 56. Dixon G, Matz M. Changes in gene body methylation do not correlate with changes in gene expression in Anthozoa or Hexapoda. BMC Genom 2022;23:234.
- 57. Feil R, Fraga MF. Epigenetics and the environment: emerging patterns and implications. Nat Rev Genet 2012;13:97-109.
- 58. Yi SV, Makova K. Insights into epigenome evolution from animal and plant methylomes. *Genome Biol Evol* 2017;9:3189–201.
- Jablonka E, Lamb MJ. The changing concept of epigenetics. Ann N Y Acad Sci 2002;981:82–96.
- 60. Weber M, Hellmann I, Stadler MB et al. Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome. Nat Genet 2007;**39**:457–66.
- 61. Sol Dourdin T, Guyomard K, Rabiller M et al. Ancestors' gift: parental early exposure to the environmentally realistic pesticide mixture drives offspring phenotype in a larger extent than direct exposure in the pacific oyster, *Crassostrea gigas*. Environ Sci Technol 2024;**58**:1865–76.

- 62. Gavery MR, Roberts SB. Predominant intragenic methylation is associated with gene expression characteristics in a bivalve mollusc. *PeerJ* 2013;**1**:e215.
- 63. Gavery M, Roberts S. Genetics & epigenetics in life history and reproduction: oysters. In: Skinner MK (ed.), *Encyclopedia of Reproduction*, 2nd edn. Oxford: Academic Press, 2018, 736–42.
- 64. Belardin LB, Antoniassi MP, Camargo M et al. Semen levels of matrix metalloproteinase (MMP) and tissue inhibitor of metallorproteinases (TIMP) protein families members in men with high and low sperm DNA fragmentation. Sci Rep 2019;9:903.
- 65. Cabral-Pacheco GA, Garza-Veloz I, Castruita-de la Rosa C *et al*. The roles of matrix metalloproteinases and their inhibitors in human diseases. Int J Mol Sci 2020;**21**:9739.
- 66. Riviere G, Fellous A, Franco A *et al*. A crucial role in fertility for the oyster angiotensin-converting enzyme orthologue CgACE. *PLoS One* 2011;**6**:e27833.
- 67. Roberts SB, Gavery MR. Is there a relationship between DNA methylation and phenotypic plasticity in invertebrates? *Front Physiol* 2012;**2**:116.
- 68. Song K, Li L, Zhang G. The association between DNA methylation and exon expression in the Pacific oyster Crassostrea gigas. PLoS One 2017;12:e0185224.
- 69. Bogan SN, Strader ME, Hofmann GE. Associations between DNA methylation and gene regulation depend on chromatin accessibility during transgenerational plasticity. BMC Biol 2023;**21**:149.
- 70. Feely RA, Alin SR, Newton J et al. The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. Estuarine Coastal Shelf Sci 2010;88:442–49.
- 71. Whitlock MC, Schluter D. The Analysis of Biological Data. London: Macmillan Higher Education, 2019.
- 72. Helm MM, Bourne N. Hatchery Culture of Bivalves: A Practical Manual. Rome: Food and Agriculture Organization of the United Nations, 2004.
- 73.Bogdanova EA, Barsova EV, Shagina IA et al. Normalization of full-length-enriched cDNA. Methods Mol Biol 2011;729: 85–98.
- 74. Gómez-Chiarri M, Warren WC, Guo X et al. Developing tools for the study of molluscan immunity: the sequencing of the genome of the eastern oyster, *Crassostrea virginica*. Fish Shellfish Immunol 2015;**46**:2–4.
- 75. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 2010;**26**: 841–42.
- 76. Tarailo-Graovac M, Chen N. Using RepeatMasker to identify repetitive elements in genomic sequences. Curr Protoc Bioinf 2009;25:4–10.
- 77. Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0 2013–2015. 2013.
- 78. White S, Roberts S. C.virginica methylation and gene expression. 2022.

- 79. Di Tommaso P, Chatzou M, Floden EW et al. Nextflow enables reproducible computational workflows. Nat Biotechnol 2017;**35**:316–19.
- 80. Nunn A, Can SN, Otto C *et al.* EpiDiverse toolkit: a pipeline suite for the analysis of bisulfite sequencing data in ecological plant epigenetics. NAR *Genom Bioinform* 2021;**3**:1qab106.
- Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for bisulfite-seq applications. *Bioinformatics* 2011;27:1571–72.
- 82. Korneliussen TS, Moltke I. NgsRelate: a software tool for estimating pairwise relatedness from next-generation sequencing data. *Bioinformatics* 2015;**31**:4009–11.
- 83. Silliman K, Spencer LH, White SJ *et al*. Epigenetic and genetic population structure is coupled in a marine invertebrate. *Genome Biol Evol* 2023;**15**:evad013.
- Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;**15**:550.
- Chen S, Zhou Y, Chen Y et al. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 2018;34:i884–90.
- 86. Ewels P. MultiQC. GitHub.
- Kim D, Paggi JM, Park C et al. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 2019;**37**:907–15.
- Pertea M, Kim D, Pertea GM et al. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat Protoc 2016;11:1650–67.
- Pertea M, Pertea GM, Antonescu CM et al. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol 2015;33:290–95.
- 90. R Core Team. R: A Language and Environment for Statistical Computing. 2021.
- 91. Fu J, Frazee AC, Collado-Torres L et al. ballgown: flexible, isoformlevel differential expression analysis. 2021.
- 92. Wickham H, Averick M, Bryan J et al. Welcome to the tidyverse. J Open Source Software 2019;**4**:1686.
- 93. Alexa A, Rahnenfuhrer J. topGO: enrichment analysis for gene ontology. R Package Version 2010;**2**:2010.
- 94. Jarmund AH, Madssen TS, Giskeødegård GF. ALASCA: an R package for longitudinal and cross-sectional analysis of multivariate data by ASCA-based methods. Front Mol Biosci 2022;9:962431.
- Pattengale ND, Alipour M, Bininda-Emonds ORP et al. How many bootstrap replicates are necessary? J Comput Biol 2010;17:337–54.
- 96. Gao S, Zou D, Mao L et al. BS-SNPer: SNP calling in bisulfite-seq data. Bioinformatics 2015;**31**:4006–08.
- 97. Akalin A, Kormaksson M, Li S et al. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol* 2012;**13**:R87.
- 98. Wu X, Lindsey ARI, Chatterjee P et al. Distinct epigenomic and transcriptomic modifications associated with Wolbachia-mediated asexuality. PLoS Pathog 2020;16:e1008397.

© The Author(s) 2024. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (https://creativecommons.org/ licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site–for further information please contact journals.permissions@oup.com.

Environmental Epigenetics, 2024, 10(1), dvae018, DOI: https://doi.org/10.1093/eep/dvae018, Advance Access Publication 23 September 2024, Research Article

Received 4 April 2024; revised 15 August 2024; accepted 22 September 2024